



Tobacco Use: Screening and Cessation Intervention

Date: 3/2/20

Description: *Percentage of patients aged 18 years and older who were screened for tobacco use one or more times within 24 months AND who received tobacco cessation intervention if identified as a tobacco user*

Initial Population: All patients aged 18 years and older seen for at least two visits or at least on preventive visit during the measurement period

Denominator Pop 1: Equals Initial Population

Denominator Pop 2: Equals Initial Population who were screened for tobacco use and identified as a tobacco user

Denominator Pop 3: Equals Initial Population

Numerator Pop 1: Patients who were screened for tobacco use at least once within 24 months


Numerator Pop 2: Patients who received tobacco cessation intervention

Numerator Pop 3: Patients who were screened for tobacco use at least once within 24 months AND who received tobacco cessation intervention if identified as a tobacco user

Exclusions: Documentation of medical reason for not screening for screening for tobacco use (limited life expectancy, other medical reason)

Allscripts Tobacco Use Screening:

Most of our patients already have a tobacco status documented due to Meaningful Use, however, the status has not been reassessed in a few years. Ask the patient their current smoking status. On the **Problem** tab, change the filter to **Social History**. If the status is correct, click on the piece of paper to assess the problem.



Allscripts Tobacco Cessation Intervention:

If you have a patient that is a tobacco user, **action is required.** Ordering a medication or providing tobacco cessation counseling will close the gap.



Tobacco Cessation

Chantix 0.5 MG Oral Tablet

Chantix 1 MG Oral Tablet

Chantix Continuing Month Pak 1 MG Oral Tablet

Chantix Starting Month Pak 0.5 MG X 11 & 1 MG X 42 Oral Tablet

Smoking and tobacco-use cessation counseling visit, 3-10 mins (Smoking and tobacco-use cessation counseling, 3 to 10 mins.)

Smoking and tobacco-use cessation counseling visit; >10min (Smoking and tobacco-use cessation counseling, >10min)